

Brane-Bulk Interaction and Holographic Principle

Yun Soo Myung*

Relativity Research Center and School of Computer Aided Science
Inje University, Gimhae 621-749, Korea

Jin Young Kim†

Department of Physics, Kunsan National University, Kunsan 573-701, Korea
Asia Pacific Center for Theoretical Physics, Pohang 790-784, Korea

Abstract

We introduce the brane-bulk interaction to discuss a limitation of the cosmological Cardy-Verlinde formula which is useful for the holographic description of brane cosmology. In the presence of the brane-bulk interaction, we cannot find the entropy representation of the first Friedmann equation (the cosmological Cardy-Verlinde formula). In the absence of the interaction, the cosmological Cardy-Verlinde formula is established even for the time-dependent charged AdS background. Hence, if there exists a dynamic exchange of energy between the brane and the bulk (that is, if $\tilde{T}^t_y \neq 0$), we cannot achieve the cosmological holographic principle on the brane.

*email address: ysmyoung@physics.inje.ac.kr

†email address: jkim@kunsan.ac.kr

Recently the brane cosmology have received much attention [1–5] in connection with the brane world scenario [6]. Further the brane cosmological evolution was explored when the brane-bulk interaction exists [7–11]. On the other hand the cosmological holographic principle was realized by applying the AdS/CFT correspondence [12–18].

In this letter we wish to clarify the relationship between the cosmological holographic principle and the brane-bulk energy exchange.

We start with reviewing the two important issues. One is a static version called the Cardy-Verlinde formula which shows that using the AdS/CFT correspondence [19], the entropy of a conformal field theory (CFT) in any dimension can be expressed in terms of a generalized form of the Cardy formula [20]. Explicitly one can consider a CFT residing in an $(n + 1)$ -dimensional spacetime with the static metric for the Einstein space $ds_{CFT}^2 = -d\tau^2 + r^2 d\Omega_n^2$, where $d\Omega_n^2$ denotes a unit n -dimensional sphere. For a strongly coupled CFT with AdS dual, one can obtain the Cardy-Verlinde formula which states that an entropy (S) can be expressed as a square root of the energy (E)

$$S = \frac{2\pi r}{n} \sqrt{E_c(2E - E_c)} \quad (1)$$

with the Casimir energy (E_c) for a finite system. Indeed, this formula was checked to hold for various kinds of AdS-bulk spacetimes; AdS-Schwarzschild black hole [12], AdS-Kerr black hole [21], AdS-charged black hole [22], AdS-Taub-Bolt spacetimes [23,24].

The other is a dynamic version called the cosmological Cardy-Verlinde formula which connects the Cardy-Verlinde formula Eq.(1) with the first Friedmann equation. We introduce the Friedmann-Robertson-Walker (FRW) metric based on the isotropic and homogeneous universe

$$ds_{FRW}^2 = -d\tau^2 + \mathcal{R}^2(\tau) d\Omega_n^2 \quad (2)$$

with a scale factor $\mathcal{R}(\tau)$. Although two line elements ds_{CFT}^2 and ds_{FRW}^2 have different origins, these are considered to be conformally equivalent. Hence it is possible to make a connection between these if some conditions are satisfied. For a closed ($k = 1$) radiation-dominated universe, two Friedmann equations are given by

$$H^2 = \frac{16\pi G_{n+1}}{n(n-1)} \rho - \frac{1}{\mathcal{R}^2}, \quad (3)$$

$$\dot{H} = -\frac{8\pi G_{n+1}}{n-1} (\rho + p) + \frac{1}{\mathcal{R}^2}, \quad (4)$$

where $H = \dot{\mathcal{R}}/\mathcal{R}$ is the Hubble parameter and the dot stands for the differentiation with respect to the proper time τ . $\rho = E/V$ is the energy density where E is the energy of radiation-matter filling the universe, $V = \mathcal{R}^n V(S^n)$ is the volume of the universe, and $p = \rho/n$ denotes the radiation-pressure. Verlinde [12] have proposed that the first Friedmann equation (3) can be expressed in terms of three cosmological holographic entropies [25,26]; Bekenstein-Verlinde entropy $S_{BV} = \frac{2\pi}{n} E\mathcal{R}$, Bekenstein-Hawking entropy $S_{BH} = (n-1) \frac{V}{4G_{n+1}\mathcal{R}}$, Hubble entropy $S_H = (n-1) \frac{HV}{4G_{n+1}}$. Then the first Friedmann equation (3) can be rewritten as an entropy relation

$$S_H^2 + (S_{BV} - S_{BH})^2 = S_{BV}^2. \quad (5)$$

Making use of the Bekenstein-Verlinde entropy as $S_{BH} \equiv \frac{2\pi}{n} E_{BH} \mathcal{R}$, we define the Bekenstein-Hawking energy $E_{BH} = \frac{n(n-1)V}{8\pi G_{n+1} \mathcal{R}^2}$. Then Eq.(5) leads to the cosmological Cardy-Verlinde formula

$$S_H = \frac{2\pi \mathcal{R}}{n} \sqrt{E_{BH}(2E - E_{BH})}. \quad (6)$$

It is straightforward to recover the Cardy-Verlinde formula of Eq.(1) from Eq.(6) replacing $(S_H, E_{BH}, \mathcal{R})$ by (S, E_c, r) respectively. This apparent connection between the static Cardy-Verlinde formula and the first Friedmann equation comes mainly from the fact that two line elements (ds_{CFT}^2 and ds_{FRW}^2) are conformally equivalent to each other. The difference is their origin that one is static, while the other is dynamic. This observation implies that the Friedmann equation and the Cardy-Verlinde formula are closely related. Furthermore it was suggested that these can be derived from the same principle.

At this stage, we ask of how much the Cardy-Verlinde formula of Eq.(1) is useful for describing a cosmological holographic principle in the brane cosmology¹. In general, one says that if the dynamic equation of the brane (the first Friedmann equation) is expressed in terms of the bulk quantities, the cosmological holographic principle is realized. In this direction, Savonije and Verlinde [27] have made the first progress by introducing a concrete model, an MDW in the $(n+2)$ -dimensional AdS-Schwarzschild black hole. There exists a remarkable relationship between the thermodynamics of a CFT on the brane and the gravitational dynamics of the brane in dual AdS space. It is well-known that the dynamic equations governing the motion of the brane are exactly given by the $(n+1)$ -dimensional Friedmann equations. Actually all moving domain walls in the AdS-Schwarzschild black hole may take a kind of holographic matter ($\rho_h = E/V, E \sim m\ell/\mathcal{R}$) which comes originally from the Schwarzschild mass-term m/r^{n-1} . One identifies this either a holographic term of strongly coupled CFT-radiation matter filling the brane or a candidate for dark matter [28]. In this work we keep a view of the holographic term. In this sense one insists that the cosmological holography can be realized by the extension of the AdS/CFT correspondence. Furthermore it was shown that the cosmological Cardy-Verlinde formula Eq.(6) matches exactly with the Cardy-Verlinde formula Eq.(1) when the brane crosses the black hole horizon. This means that we can achieve the equalities $S = S_H, E_c = E_{BH}$ at a holographic point $r = r_H = \mathcal{R}_H$. This is an important result in the sense that from the Friedmann equations we know about thermodynamics of the CFT-radiation matter filled in the universe. The second Friedmann equation (4) leads to the Smarr-type formula of $E_{BH} = n(E + pV - \mathcal{T}_H S_H)$ with the Hubble temperature defined by $\mathcal{T}_H = -\frac{\dot{H}}{2\pi H}$. It is related to the defining equation of the Casimir energy on the CFT side: $E_c = n(E + pV - TS)$.

In the MDW approach the bulk spacetime is fixed. This means that we can derive the gravitational dynamics of the brane from the given bulk black hole background. However, we don't know precisely what kinds of higher dimensional theories we should start with, even if string theories provide us some informations about it. In this sense the MDW approach is considered as a restricted one although it provides us an exact evolution of the brane.

¹There exist two approaches for the brane cosmology: one is the moving domain wall (MDW) approach [4,5] and the other is Binetruy-Deffayet-Langlois (BDL) approach [1,2].

On the other hand the BDL approach is a genuine extension of the Kaluza-Klein cosmology to account for the local distribution on the brane. In this case the location of the brane is fixed with respect to the bulk direction. Also this approach is useful for describing the cosmological evolution of the brane when a brane-bulk interaction exists. But there are some ambiguities in interpreting the holographic matter. From now on, we follow the BDL brane cosmology [1,2]. We introduce the Gaussian-normal bulk metric for $(1 + 3 + 1)$ -dimensional spacetime

$$ds_{BDL}^2 = -c^2(t, y)dt^2 + a^2(t, y)\gamma_{ij}dx^i dx^j + b^2(t, y)dy^2, \quad (7)$$

where γ_{ij} is the metric of a three-dimensional space with a constant curvature of $6k$. Let us express the bulk Einstein equation $G_{MN} = \frac{1}{2M^3}T_{MN}$ in terms of the BDL metric ². Now we introduce a $(1 + 3)$ -dimensional brane located at $y = 0$. We choose for simplicity the same stress-energy tensor $T^M{}_N = \text{diag}(-\Lambda, -\Lambda, -\Lambda, -\Lambda, -\Lambda) + \tilde{T}^M{}_N$ on both sides. Here the bulk cosmological constant is given by Λ . The bulk stress-energy tensor $\tilde{T}^M{}_N$ from $\tilde{\mathcal{L}}_B^{mat}$ is not needed to have a specific form initially. If $\tilde{T}^t{}_y = 0$, it is obvious that there is no brane-bulk interaction in a dynamical way. The local stress-energy tensor from $\mathcal{L}_b^{mat}$ including the brane tension σ is assumed to be of the form

$$\tau^\mu{}_\nu = \frac{\delta(y)}{b} \text{diag}(-\rho - \sigma, p - \sigma, p - \sigma, p - \sigma, 0). \quad (8)$$

We are interested in solving the Einstein equations at the location of the brane. We indicate by the subscript “0” the value of the various quantities on the brane. Also it is convenient to choose the Gaussian-normal gauge with $b_0 = 1$ and the temporal gauge with $c_0 = 1$ on the brane. We obtain, from $G_{0y} = \frac{1}{2M^3}T_{0y}$,

$$\dot{\rho} + 3\frac{\dot{a}_0}{a_0}\rho(1 + \omega) = -2\tilde{T}^0{}_y. \quad (9)$$

Here we assumed an equation of state $p = \omega\rho$. On the other hand, the average part of yy -component equation is given by

$$\frac{\ddot{a}_0}{a_0} + \left(\frac{\dot{a}_0}{a_0}\right)^2 + \frac{k}{a_0^2} = \frac{1}{6M^3}\left(\Lambda + \frac{\sigma^2}{12M^3}\right) - \frac{1}{144M^6}\left(\sigma(3p - \rho) + \rho(3p + \rho)\right) - \frac{1}{6M^3}\tilde{T}^y{}_y. \quad (10)$$

Then we rewrite Eq.(10) in the following equivalent form

$$H_0^2 = \frac{1}{144M^6}(\rho^2 + 2\sigma\rho) - \frac{k}{a_0^2} + \chi + \phi + \frac{1}{12M^3}\left(\Lambda + \frac{\sigma^2}{12M^3}\right), \quad (11)$$

$$\dot{\chi} + 4H_0\chi = \frac{1}{36M^6}(\rho + \sigma)\tilde{T}^0{}_y, \quad (12)$$

$$\dot{\phi} + 4H_0\phi = -\frac{1}{3M^3}H_0\tilde{T}^y{}_y, \quad (13)$$

²Our action is given by $S_5 = \int d^5x \sqrt{-g}(M^3R - \Lambda + \tilde{\mathcal{L}}_B^{mat}) + \int d^4x \sqrt{-\hat{g}}\mathcal{L}_b^{mat}$ with $M^3 = 1/16\pi G_5$, $\mathcal{L}_b^{mat} = -\sigma + \dots$.

with $H_0 = \dot{a}_0/a_0$. In the case of $p = \rho = 0$ and $\phi = \chi = 0$, we wish to find the Randall-Sundrum vacuum [6]. Hence we choose the bulk cosmological constant $\Lambda = -\sigma^2/12M^3 = -12M^3/\ell^2$ and the brane tension $\sigma = 12M^3/\ell$ to have a critical brane. Hence the cosmological evolution is determined by four initial parameters $(\rho_i, a_{0i}, \chi_i, \phi_i)$ instead of the two (ρ_i, a_{0i}) . This is so because the generalized Friedmann equation (11) is not a first integral of the Einstein equations. It is mainly due to the energy exchange $\tilde{T}^t_y$ between the brane and the bulk. In the case of $\phi = 0, \tilde{T}^t_y = A\rho > 0$ and negligible $\tilde{T}^y_y$, one finds a mirage-radiation term $\chi \sim (1 - e^{-At/2})/a_0^4$ for an energy outflow from the brane [10]. It can be considered that the real brane matter decays into the extra dimension. Also for $\phi = 0, \tilde{T}^t_y \sim -\frac{1}{a_0^q}$ and negligible $\tilde{T}^y_y$, it is shown that the energy influx from the bulk generates a cosmological acceleration on the brane with the acceleration parameter $Q \equiv \frac{1}{H_0^2} \frac{\ddot{a}_0}{a_0} = 1 - \frac{q}{3}$ where $0 \leq q \leq 4$ [11]. However, in general it will be a formidable task to obtain the solution of Eqs.(9)-(13) because it gives rise to a complicated dynamics between the brane and the bulk. Importantly it is very hard to express Eq.(11) in terms of the cosmological Cardy-Verlinde formula of Eq.(6) because Eq.(11) is not a first integral of the Einstein equations.

In this work we are interested in exploring the relationship between the brane-bulk interaction and the (cosmological) holographic principle. In the case of $\tilde{T}^t_y = \tilde{T}^y_y = 0$, it is shown from Eqs.(12)-(13) that one finds the mirage or Weyl-radiation terms of $\chi = C_\chi/a_0^4$ and $\phi = C_\phi/a_0^4$ with two unknown constants C_χ, C_ϕ . In view of the moving domain wall approach these correspond to the holographic-radiation terms originated from the mass of the AdS-black hole [29]. Furthermore, it is obvious that the omission of ϕ is not justified when the energy influx (outflux) stops and thus $\tilde{T}^t_y$ becomes zero. The authors in [10,11] assumed that at this moment, $\tilde{T}^y_y$ is also very small, so that ϕ gives a negligible contribution to Eq.(11). However, this is not the general case to obtain the information on the brane.

We propose here that *as far as there exists any dynamical interaction between the brane and the bulk, one cannot establish the cosmological holographic principle*. The reason is as follows. First, up to now one does not find any dynamical localization mechanism for the brane matter by choosing an appropriate interaction term $\tilde{T}^t_y$. This means that we do not specify the details of the mechanism through which energy is transferred from the bulk to the brane. Second the holographic principle is based on the static non-local mechanism, for example, the projection of the bulk Weyl tensor onto the brane [30]. The AdS/CFT correspondence is a tool to realize the holographic principle. In the case of brane cosmology, we obtain the holographic energy density ρ_h on the brane by the extension of the AdS/CFT correspondence. Actually, one does not want to have an environment where a dynamical energy exchange between the brane and bulk exists. This is so because if $\tilde{T}^t_y \neq 0$, one cannot identify which part belongs to the brane and which part belongs to the bulk. Finally, if $\tilde{T}^t_y = 0$, Eq.(11) becomes a first integral (that is, the first Friedmann equation). As a result, to establish the holographic principle, one has to meet the condition of no dynamical exchange between the brane and bulk : $\tilde{T}^t_y = 0$.

In order to see this with a specific example, let us introduce the bulk Maxwell field whose solution takes the form under the metric of Eq.(7) [31,32]

$$F_{yt} = \frac{\mathcal{Q}bc}{a^n}, \quad (14)$$

where $\mathcal{Q}$ is an unknown integration constant which is proportional to the charge of the

AdS-charged black hole if one makes a further connection to the MDW approach. Then the bulk stress-energy tensor $\tilde{T}^M{}_N = 4M^3(F^{MP}F_{NP} - \delta^M{}_N F^2/4)$ is given by³

$$\tilde{T}^M{}_N = 2M^3 \text{diag} \left(-\frac{\mathcal{Q}^2}{a^6}, +\frac{\mathcal{Q}^2}{a^6}, \dots, -\frac{\mathcal{Q}^2}{a^6} \right). \quad (15)$$

This implies that on the brane, $\tilde{T}^t{}_y = 0, \tilde{T}^y{}_y = -2M^3\mathcal{Q}^2/a_0^6$. Although this is a time-dependent bulk term, there does not exist any dynamical exchange between the brane and bulk. Substituting these into Eqs.(12)-(13), one finds that

$$\phi = -\mathcal{Q}^2/3a_0^6, \quad \chi = \mathcal{C}_\chi/a_0^4. \quad (16)$$

Equation (11) leads to the first Friedmann equation

$$H_0^2 = -\frac{k}{a_0^2} + \frac{\mathcal{C}_\chi}{a_0^4} - \frac{1}{3} \frac{\mathcal{Q}^2}{a_0^6} + \frac{1}{144M^6}(\rho^2 + 2\sigma\rho). \quad (17)$$

This is an equation which is derived from the BDL approach and governs the evolution of the fixed brane sandwiched in the charged AdS bulks. For $\rho = p = 0$ case, let us compare our equation (17) with that of the moving domain wall in the charged topological-AdS (CTAdS) black holes which can be rewritten as [33–35]

$$H^2 = -\frac{k}{\mathcal{R}^2} + \frac{\omega_3 M}{\mathcal{R}^4} - \frac{3\omega_3^2 Q^2}{16\mathcal{R}^6}, \quad (18)$$

where $\omega_3 = 1/3M^3V(S^3)$. This can be written as the cosmological Cardy-Verlinde formula as

$$S_H = \frac{2\pi\mathcal{R}}{3} \sqrt{E_{BH}[2(E - \Phi Q/2) - kE_{BH}]}, \quad (19)$$

with the electric potential $\Phi = 3\ell\omega_3 Q/16\mathcal{R}^3$ on the brane. Further its time-derivative is given by

$$kE_{BH} = 3(E + pV - \Phi Q - \mathcal{T}_H S_H). \quad (20)$$

With the identification,

$$a_0 \leftrightarrow \mathcal{R}, \quad \mathcal{C}_\chi \leftrightarrow \omega_3 M, \quad \mathcal{Q} \leftrightarrow \pm \frac{3\omega_3 Q}{4} \quad (21)$$

our equation (17) with $p = \rho = 0$ coincides with the cosmological Cardy-Verlinde formula Eq.(19) in the presence of the background charge. This shows that the cosmological holographic principle is achieved.

In case of Eq. (18) for the moving brane, the bulk is just fixed as the two CTAdS black holes with the mass M and electric charge Q . These two parameters encode information of

³To obtain this, one takes $\tilde{\mathcal{L}}_B^{mat} = -M^3 F^2$ in footnote 2.

the bulk and are used to describe a strongly coupled CFT on the brane via the extension of the AdS/CFT correspondence. On the other hand, in deriving Eq. (17), we don't need to know about the precise form of the AdS bulk geometry but we have to keep up a bulk Maxwell field. However, the two integration constants $\mathcal{C}_\chi$ and $\mathcal{Q}$ certainly encode information of the bulk, as the initial conditions. Just as in the moving domain wall approach in which the brane moves in the fixed bulk and acts as the boundary of the bulk, the Friedmann equation of the BDL brane sandwiched in two AdS bulks carries the information of the bulk as the initial conditions. This shows that it is possible to achieve the cosmological holographic principle in the BDL approach. Furthermore, to make a further connection with the static Cardy-Verlinde formula, we have to choose the CTAdS black hole background as the fixed bulk [31,32].

In conclusion, if there exists a dynamic exchange of energy between the brane and the bulk (that is, if $\tilde{T}^t{}_y \neq 0$), one cannot achieve the cosmological holographic principle on the brane.

ACKNOWLEDGEMENTS

This work was supported in part by KOSEF, Project Number R02-2002-000-00028-0.

REFERENCES

- [1] P. Binetruiy, C. Deffayet and D. Langlois, Nucl. Phys. **B565**, 269 (2000) [hep-th/9905012].
- [2] P. Binetruiy, C. Deffayet, U. Ellwanger and D. Langlois, Phys. Lett. **B477**, 285 (2000) [hep-th/9910219].
- [3] H. Chamblin and H. Real, Nucl. Phys. B **562**, 133 (1999) [hep-th/9903225].
- [4] P. Kraus, JHEP **9912**, 011 (1999) [hep-th/9910149].
- [5] D. Ida, JHEP **0009**, 014 (2000) [gr-qc/9912002].
- [6] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 4690 (1999) [hep-th/9906064].
- [7] C. van de Bruck, M. Dorca, C. J. A. P. Martins, and M. Parry, Phys. Lett. B **495**, 183 (2000) [hep-th/0009056].
- [8] A. Hebecker and J. March-Russell, Nucl. Phys. B **608**, 375(2001) [hep-th/0103214].
- [9] D. Langlios, L. Sorbo and M. Rodriguez-Martinez, Phys. Rev. Lett. **89**, 171301(2002) [hep-th/0206146].
- [10] E. Kiritsis, G. Kofinas, N. Tetradis, T. N. Tomaras and V. Zarikas, JHEP **0302**, 035(2003) [hep-th/0207060].
- [11] N. Tetradis, “Cosmological Acceleration from Energy Influx”, hep-th/0211200.
- [12] E. Verlinde, “On the Holographic Principle in a Radiation Dominated Universe”, hep-th/0008140.
- [13] F.-L. Lin, Phys. Rev. D **63**, 064026 (2001) [hep-th/0010127].
- [14] S. Nojiri and S. Odintsov, Int. J. Mod. Phys. A **16**, 3273 (2001) [hep-th/0011115]; Class. Quant. Grav. **18**, 5227(2001) [hep-th/0103078].
- [15] B. Wang, E. Abdalla and R.K. Su, Phys. Lett. B **503**, 394(2001) [hep-th/0101073].
- [16] R. Brustein, S. Foffa and G. Veneziano, Phys. Lett. B **507**, 270 (2001) [hep-th/0101083].
- [17] D. Klemm, A.C. Petkou, G. Siopsis and D. Zanon, Nucl. Phys. B **620**, 519 (2002) [hep-th/0104141].
- [18] D. Youm, Mod. Phys. Lett. A **16**, 1236 (2001) [hep-th/0105036].
- [19] J. Maldacena, Adv. Theor. Math. Phys. **2**, 231 (1998) [hep-th/9711200]; S. S. Gubser, I. Klebanov, A. Polyakov, Phys. Lett. B **428**, 105 (1998) [hep-th/9802109]; E. Witten, Adv. Theor. Math. Phys. **2**, 253 (1998) [hep-th/9802150].
- [20] J.L. Cardy, Nucl. Phys. **B270**, 186 (1986).
- [21] D. Klemm, A.C. Petkou and G. Siopsis, Nucl. Phys. B **601**, 380 (2001) [hep-th/0101076].
- [22] R.G. Cai, Phys. Rev. D **63**, 124018 (2001) [hep-th/0102113].
- [23] D. Birmingham and S. Mokhtari, Phys. Lett. B **508**, 365 (2001) [hep-th/0103108].
- [24] E. Witten, Adv. Theor. Math. Phys. **2**, 505 (1998) [hep-th/9803131].
- [25] J.D. Bekenstein, Phys. Rev. D **23**, 287 (1981).
- [26] R. Bousso, JHEP **0004**, 035 (2001) [hep-th/0012052]; JHEP **0011**, 038 (2000) [hep-th/0010252].
- [27] I. Savonije and E. Verlinde, Phys. Lett. B **507**, 305 (2001) [hep-th/0102042].
- [28] A. Coley, Phys. Rev. D **66**, 023512 (2002) [hep-th/0110049].
- [29] N.J. Kim, H.W. Lee, Y.S. Myung and G. Kang, Phys. Rev. D **64**, 064022 (2001) [hep-th/0104159].
- [30] T. Shiromizu, K. Maeda, and M. Sasaki, Phys. Rev. D **62**, 024012 (2000) [gr-qc/9910076].

- [31] R.G. Cai, Y. S. Myung and N. Ohta, Class. Quant. Grav. **18**, 5429 (2001) [hep-th/0105070].
- [32] Y. S. Myung, Class. Quant. Grav. **20**, 935 (2003) [hep-th/0208086].
- [33] A.K. Biswas and S. Mukherji, JHEP **0103**, 046 (2001) [hep-th/0102138].
- [34] Y. S. Myung, Mod. Phys. Lett. A **17**, 1915(2002) [hep-th/0103241].
- [35] P. Kanti and K. Tamvakis, “Challenges and Obstacles for a Bouncing Universe in Brane Models”, hep-th/0303073.